


Listing of Claims:

Please rewrite claims 49, 59, 69 and 79 and cancel claims 50-53, 60-63, 70-73 and 80-83 as follows.

1-48. (Canceled)

49. (Currently Amended) A method of treating a cancerous disease selected from the group consisting of leukemia, colon cancer, breast cancer and prostate cancer comprising administering to a patient with said disease an effective amount of 20(S)-1 α ,25-dihydroxy-2-methylene-26,27-dihomo-19-norvitamin D₃ having the structure



50-53. (Canceled)

54. (Previously Presented) The method of claim 49 wherein the compound is administered orally.


55. (Previously Presented) The method of claim 49 wherein the compound is administered parenterally.

56. (Previously Presented) The method of claim 49 wherein the compound is administered transdermally.

57. (Previously Presented) The method of claim 49 wherein said effective amount comprises about 0.01 μ g/day to about 100 μ g/day of 20(S)-1 α ,25-dihydroxy-2-methylene-26,27-dihomo-19-norvitamin D₃.

58. (Previously Presented) The method of claim 49 wherein said effective amount comprises about 0.1 μ g/day to about 50 μ g/day of 20(S)-1 α ,25-dihydroxy-2-methylene-26,27-dihomo-19-norvitamin D₃.

59. (Currently Amended) A method of treating a cancerous disease selected from the group consisting of leukemia, colon cancer, breast cancer and prostate cancer comprising administering to a patient with said disease an effective amount of 20(S)-26,27-dimethylene-25-methoxy-2-methylene-19-norvitamin D₃ having the structure



60-63. (Canceled)

64. (Previously Presented) The method of claim 59 wherein the compound is administered orally.


65. (Previously Presented) The method of claim 59 wherein the compound is administered parenterally.

66. (Previously Presented) The method of claim 59 wherein the compound is administered transdermally.

67. (Previously Presented) The method of claim 59 wherein said effective amount comprises about 0.01 μ g/day to about 100 μ g/day of 20(S)-26,27-dimethylene-25-methoxy-2-methylene-19-norvitamin D₃.

68. (Previously Presented) The method of claim 59 wherein said effective amount comprises about 0.1 $\mu\text{g/day}$ to about 50 $\mu\text{g/day}$ of 20(S)-26,27-dimethylene-25-methoxy-2-methylene-19-norvitamin D₃.

69. (Currently Amended) A method of treating a cancerous disease selected from the group consisting of leukemia, colon cancer, breast cancer and prostate cancer comprising administering to a patient with said disease an effective amount of 20(S)-1 α ,25-dihydroxy-26,27-dimethylene-2-methylene-19-norvitamin D₃ having the structure



70-73. (Canceled)

74. (Previously Presented) The method of claim 69 wherein the compound is administered orally.


75. (Previously Presented) The method of claim 69 wherein the compound is administered parenterally.

76. (Previously Presented) The method of claim 69 wherein the compound is administered transdermally.

77. (Previously Presented) The method of claim 69 wherein said effective amount comprises about 0.01 $\mu\text{g/day}$ to about 100 $\mu\text{g/day}$ of 20(S)-1 α ,25-dihydroxy-26,27-dimethylene-2-methylene-19-norvitamin D₃.

78. (Previously Presented) The method of claim 69 wherein said effective amount comprises about 0.1 $\mu\text{g/day}$ to about 50 $\mu\text{g/day}$ of 20(S)-1 α ,25-dihydroxy-26,27-dimethylene-2-methylene-19-norvitamin D₃.

79. (Currently Amended) A method of treating a cancerous disease selected from the group consisting of leukemia, colon cancer, breast cancer and prostate cancer comprising administering to a patient with said disease an effective amount of 20(S)-26,27-dimethylene-1 α -hydroxy-2-methylene-24-dehydro-19-norvitamin D₃ having the structure



80-83. (Canceled)

84. (Previously Presented) The method of claim 79 wherein the compound is administered orally.

85. (Previously Presented) The method of claim 79 wherein the compound is administered parenterally.

86. (Previously Presented) The method of claim 79 wherein the compound is administered transdermally.

87. (Previously Presented) The method of claim 79 wherein said effective amount comprises about 0.01 μ g/day to about 100 μ g/day of 20(S)-26,27-dimethylene-1 α -hydroxy-2-methylene-24-dehydro-19-norvitamin D₃.

88. (Previously Presented) The method of claim 79 wherein said effective amount comprises about 0.1 μ g/day to about 50 μ g/day of 20(S)-26,27-dimethyl-1 α -hydroxy-2-methylene-24-dehydro-19-norvitamin D₃.